

Walk through Combinatorics: homework #3*


~~Due 19 October 2012~~

Due 22 October 2012

Collaboration and use of external sources are permitted, but discouraged, and must be fully acknowledged and cited. Collaboration may involve only discussion; all the writing must be done individually.

The number of points for each problem is specified in brackets. The problems appear in no special order.

1. [2+1] An *incomplete binary tree* is a rooted tree in which each node is either a terminal node, or a branch. A branch node might have either one child (left or right) or both left and right children. A *descending labelling* on a tree is an assignment of labels 1 through n to the nodes, where n is the number of nodes, such that on every path from the root the labels are descending. A *descending incomplete binary tree* is an incomplete binary tree with a descending labelling on it.



Three examples of descending incomplete binary trees

*This homework is from <http://www.borisbukh.org/DiscreteMath12/hw3.pdf>.

- (a) Use generating functions to compute the number of descending incomplete binary trees on n nodes.
 - (b) Give a bijective proof of the closed formula that you found in the part (a).
2. [2+1] Let F_n be the number of functions $[n] \rightarrow \mathbb{N}$ whose image is $[r]$ for some r .
 - (a) Find the exponential generating function for F_n . (Hint: consider individual values of r .)
 - (b) Find an asymptotics for F_n . (Your answer should be of the form $F_n = A_n(1 + O(c^n))$ for some explicit A_n and an explicit value of $c < 1$.)
3. [2] Let $A \subset \mathbb{N}$ be an infinite set of integers. Let R_n be the number of ways to write n as $n = a_1 + a_2$ with $a_1, a_2 \in A$, where the order of summands does not matter (i.e. $5 = 3 + 2$ and $5 = 2 + 3$ are the same). Show that $R_n = C + o(1)$ cannot hold for all large values of n .
4. [2+(2 extra credit)] Let \mathcal{S}_n be the set of all sequences of length $2n$ with n terms that are $+1$ and n terms that are -1 , and all of whose partial sums are nonnegative. Let $C_{n,k}$ be the number sequences in \mathcal{S}_n with exactly k zero partial sums. For example, the only sequence counted by $C_{2,2}$ is $+1, +1, -1, -1$.
 - (a) Compute $F(z, w) = \sum C_{n,k} z^n w^k$ in closed form.
 - (b) (Extra credit) Suppose we pick a sequence uniformly at random from \mathcal{S}_n . What is the expectation of the number of times a partial sum of coefficients vanishes? (You may want to use a symbolic algebra system for some computations.)