


Walk through Combinatorics: homework #6*

Due 3 December 2014, at start of class

Collaboration and use of external sources are **forbidden**. If unsure, ask me first.

The number of points for each problem is specified in brackets. The problems appear in no special order.

1. [2+1] An *incomplete binary tree* is a rooted tree in which each node is either a terminal node, or a branch. A branch node might have either one child (left or right) or both left and right children. A *descending* labelling on a tree is an assignment of labels 1 through n to the nodes, where n is the number of nodes, such that on every path from the root the labels are descending. A *descending incomplete binary tree* is an incomplete binary tree with a descending labelling on it.



Three examples of descending incomplete binary trees

- (a) Use generating functions to compute the number of descending incomplete binary trees on n nodes.
 - (b) Give a bijective proof of the closed formula that you found in the part (a).
2. [2+1] Let F_n be the number of functions $[n] \rightarrow \mathbb{N}$ whose image is $[r]$ for some r .

*This homework is from <http://www.borisbukh.org/DiscreteMath14/hw6.pdf>.

- (a) Find the exponential generating function for F_n . (Hint: consider individual values of r .)
- (b) Find an asymptotics for F_n . (Your answer should be of the form $F_n = A_n(1 + O(c^n))$ for some explicit A_n and an explicit value of $c < 1$.)
3. [2] Suppose $\{a_n\}$ is a sequence that satisfies a linear recurrence, i.e., a relation of the form $a_n = \sum_{k=1}^d \gamma_k a_{n-k}$ for some constants $\gamma_1, \dots, \gamma_d$. Let $b_n = a_{3n}$. Prove that b_n also satisfies a linear recurrence.
4. [2+(2 extra credit)] Let \mathcal{S}_n be the set of all sequences of length $2n$ with n terms that are $+1$ and n terms that are -1 , and all of whose partial sums are nonnegative. Let $C_{n,k}$ be the number sequences in \mathcal{S}_n with exactly k zero partial sums. For example, the only sequence counted by $C_{2,2}$ is $+1, +1, -1, -1$.
- (a) Compute $F(z, w) = \sum C_{n,k} z^n w^k$ in closed form.
- (b) (Extra credit) Suppose we pick a sequence uniformly at random from \mathcal{S}_n . What is the expectation of the number of times a partial sum of coefficients vanishes? (You may want to use a symbolic algebra system for some computations.)