

Walk through Combinatorics: homework #2*

Due 21 September 2018, at start of class

Collaboration and use of external sources are permitted, but must be fully acknowledged and cited. For your own learning, you are advised to work individually. Collaboration may involve only discussion; all the writing must be done individually.

Homework must be submitted in L^AT_EX via e-mail. I want both the L^AT_EX file and the resulting PDF. The files must be of the form `lastname_discr_hwnum.tex` and `lastname_discr_hwnum.pdf` respectively. Pictures do not have to be typeset; a legible photograph of a hand-drawn picture is acceptable.

You can earn 0.5 points of extra credit by submitting exactly 4 problems before the start of class on 14 September 2018. Resubmissions void the extra credit.

- [1] Show that if \mathbb{N} is colored into finitely many colors, then there are three *distinct* natural numbers x, y, z of the same color such that $x + y = z$.
- Let $R(k_1, k_2)$ be the usual two-color Ramsey number.
 - [2] Show that $R(5, k) \geq \frac{1}{100}(k/\log k)^2$ using a probabilistic argument. [Hint: color edges using a biased coin. The inequality $1 - p \leq \exp(-p)$ is likely to be useful.]
 - [0; do not turn in] Rewrite the proof above as a counting argument.
- [2] Let $k > r$ be natural numbers. Consider an underdetermined system of homogeneous linear equations

$$\begin{aligned}a_{11}x_1 + a_{12}x_2 + \cdots + a_{1k}x_k &= 0, \\a_{21}x_1 + a_{22}x_2 + \cdots + a_{2k}x_k &= 0, \\&\vdots \\a_{r1}x_1 + a_{r2}x_2 + \cdots + a_{rk}x_k &= 0\end{aligned}$$

with coefficients a_{ij} that are integers and satisfy $|a_{ij}| \leq N$ for all i and j . Show that there is a solution such that all the unknowns x_1, \dots, x_k are integers that are not all zero and that satisfy $|x_j| \leq C_{k,r}N^{r/(k-r)}$. Here $C_{k,r}$ is a constant that

*This homework is from <http://www.borisbukh.org/DiscreteMath18/hw2.pdf>.

depends only on k and r . You do not have to find the smallest $C_{k,r}$ for which this holds.

4. [1] Show that for every k there is an n such that whenever subsets of $[n]$ are colored into k colors there are non-empty disjoint sets A, B, C such that the colors of six sets $A, B, C, A \cup B, A \cup C, B \cup C$ are all the same.
5. [2] Product of graphs G_1, \dots, G_n is the graph whose vertices are ordered tuples (v_1, \dots, v_n) s.t. $v_i \in G_i$, and the pair $(a_1, \dots, a_n), (b_1, \dots, b_n)$ forms an edge if and only if for each i either $a_i = b_i$ or $a_i b_i \in E(G_i)$.

Let $\alpha(G)$ denote the size of the largest independent set in G . Prove that the Ramsey number $R(\underbrace{3, \dots, 3}_n)$ is equal to $1 + \max \alpha(G_1 \times \dots \times G_n)$ where the maximum is over all graphs G_1, \dots, G_n such that $\alpha(G) = 2$.

6. [1] Show that every graph G on $n \geq 16$ vertices contains two disjoint subsets A and B of size $|A| = |B| \geq \frac{1}{4} \log_2 n$ such that induced subgraphs $G|_A$ and $G|_B$ are isomorphic.
7. For a set A of integers, put $\Sigma(A) \stackrel{\text{def}}{=} \sum_{a \in A} a$. Let $S \subset [n]$ be a set of m integers.
- (a) [2] Show that if $m \geq \log_2 n + \log_2 \log_2 n + 1$ and n is sufficiently large, then S contains two non-empty disjoint subsets A_1, A_2 such that $\Sigma(A_1) = \Sigma(A_2)$.
- (b) (Open problem; much extra credit) Prove or disprove that there is a constant C such that if $m \geq C \log_2 n$, then S contains three non-empty disjoint subsets A_1, A_2, A_3 such that $\Sigma(A_1) = \Sigma(A_2) = \Sigma(A_3)$.
8. [1+1+1] Let $n \geq 100$.

- (a) Show that in every 2-coloring of edges of K_n there are at least $n/100$ vertex-disjoint monochromatic K_3 's.
- (b) Show that in every 2-coloring of edges of K_n there are at least $n^2/1000$ edge-disjoint monochromatic K_3 's.
- (c) Show that, if P is an n -point set in \mathbb{R}^2 in general position, then P contains $n/100$ convex quadrilaterals whose convex hulls are disjoint. ('General position' here means that no line contains three or more points of P .)