

# Discrete Math: homework #2\*

## Due 22 September 2021, at 9:00am

Collaboration and use of external sources are permitted, but must be fully acknowledged and cited. For your own learning, you are advised to work individually. Collaboration may involve only discussion; all the writing must be done individually.

Homework must be submitted in L<sup>A</sup>T<sub>E</sub>X via e-mail. I want both the L<sup>A</sup>T<sub>E</sub>X file and the resulting PDF. The files must be of the form `lastname_discr_hwnum.tex` and `lastname_discr_hwnum.pdf` respectively. Pictures do not have to be typeset; a legible photograph of a hand-drawn picture is acceptable.

You can earn 1 point of extra credit by submitting exactly 4 problems before 9am on 15 September 2021. Resubmissions void the extra credit.

- [1] Show that if  $\mathbb{N}$  is colored into finitely many colors, then there are three *distinct* natural numbers  $x, y, z$  of the same color such that  $x + y = z$ .
- Let  $R(k_1, k_2)$  be the usual two-color Ramsey number.
  - [2] Show that  $R(5, k) \geq \frac{1}{100}(k/\log k)^2$  using a probabilistic argument. [Hint: color edges using a biased coin. The inequality  $1 - p \leq \exp(-p)$  is likely to be useful.]
  - [0; do not turn in] Rewrite the proof above as a counting argument.
- [2] Let  $k > r$  be natural numbers. Consider an underdetermined system of homogeneous linear equations

$$\begin{aligned}a_{11}x_1 + a_{12}x_2 + \cdots + a_{1k}x_k &= 0, \\a_{21}x_1 + a_{22}x_2 + \cdots + a_{2k}x_k &= 0, \\&\vdots \\a_{r1}x_1 + a_{r2}x_2 + \cdots + a_{rk}x_k &= 0\end{aligned}$$

with coefficients  $a_{ij}$  that are integers and satisfy  $|a_{ij}| \leq N$  for all  $i$  and  $j$ . Show that there is a solution such that all the unknowns  $x_1, \dots, x_k$  are integers that are not all zero and that satisfy  $|x_j| \leq C_{k,r}N^{r/(k-r)}$ . Here  $C_{k,r}$  is a constant that

---

\*This homework is from <http://www.borisbukh.org/DiscreteMath21/hw2.pdf>.

depends only on  $k$  and  $r$ . You do not have to find the smallest  $C_{k,r}$  for which this holds.

4. [1] Show that for every  $k$  there is an  $n$  such that whenever subsets of  $[n]$  are colored into  $k$  colors there are non-empty disjoint sets  $A, B, C$  such that the colors of six sets  $A, B, C, A \cup B, A \cup C, B \cup C$  are all the same.
5. [2] Product of graphs  $G_1, \dots, G_n$  is the graph whose vertices are ordered tuples  $(v_1, \dots, v_n)$  s.t.  $v_i \in G_i$ , and the pair  $(a_1, \dots, a_n), (b_1, \dots, b_n)$  forms an edge if and only if for each  $i$  either  $a_i = b_i$  or  $a_i b_i \in E(G_i)$ .

Let  $\alpha(G)$  denote the size of the largest independent set in  $G$ . Prove that the Ramsey number  $R(\underbrace{3, \dots, 3}_n)$  is equal to  $1 + \max \alpha(G_1 \times \dots \times G_n)$  where the maximum is over all graphs  $G_1, \dots, G_n$  such that  $\alpha(G) = 2$ .

6. [1] Show that every graph  $G$  on  $n \geq 16$  vertices contains two disjoint subsets  $A$  and  $B$  of size  $|A| = |B| \geq \frac{1}{4} \log_2 n$  such that induced subgraphs  $G|_A$  and  $G|_B$  are isomorphic.
7. For a set  $A$  of integers, put  $\Sigma(A) \stackrel{\text{def}}{=} \sum_{a \in A} a$ . Let  $S \subset [n]$  be a set of  $m$  integers.
- (a) [2] Show that if  $m \geq \log_2 n + \log_2 \log_2 n + 1$  and  $n$  is sufficiently large, then  $S$  contains two non-empty disjoint subsets  $A_1, A_2$  such that  $\Sigma(A_1) = \Sigma(A_2)$ .
- (b) (Open problem; much extra credit) Prove or disprove that there is a constant  $C$  such that if  $m \geq C \log_2 n$ , then  $S$  contains three non-empty disjoint subsets  $A_1, A_2, A_3$  such that  $\Sigma(A_1) = \Sigma(A_2) = \Sigma(A_3)$ .
8. [2+(extra credit) 2] Let  $X$  be a set. A  $n$ -letter word over alphabet  $X$  is simply an element of  $X^n$ , i.e., a sequence of  $n$  elements from  $X$ . Consider a word  $w \in ([k] \cup \{*\})^n$ . Let  $C(w)$  be the set of all the words in  $[k]^n$  that can be obtained by replacing stars by elements of  $[k]$ . For example, if  $k = 2$  then

$$C(1 * 21*) = \{11211, 11212, 12211, 12212\}.$$

In general, if  $w$  has  $m$  stars, then  $|C(w)| = 2^m$ .

- (a) Use Ramsey's theorem (or anything else) to show that there is a function  $n_0(r, m)$  such that if  $n \geq n_0(r, m)$  and  $\chi: [2]^n \rightarrow [r]$  is a coloring of  $n$ -letter words over 2-letter alphabet, then there is a word  $w \in ([2] \cup \{*\})^n$  with  $m$  stars such that for  $w' \in C(w)$  the color  $\chi(w')$  depends only on the number of occurrences of 1's and 2's in  $w'$ .

- (b) Show that the analogous statement for 3-letter alphabet is false. Namely, show that there is an  $r$  and  $m$  such that for every  $n$  there is a coloring  $\chi: [3]^n \rightarrow [r]$  so that for no  $w \in ([3] \cup \{*\})^n$  with  $m$  stars the color of a word in  $C(w)$  depends only on the number of occurrences of 1's, 2's and 3's.