

Discrete Math: homework #4*

Due 20 October 2021, at 9:00am

Collaboration and use of external sources are permitted, but must be fully acknowledged and cited. For your own learning, you are advised to work individually. Collaboration may involve only discussion; all the writing must be done individually.

Homework must be submitted in L^AT_EX via e-mail. I want both the L^AT_EX file and the resulting PDF. The files must be of the form `andrewid_discr_hwnum.tex` and `andrewid_discr_hwnum.pdf` respectively. Pictures do not have to be typeset; a legible photograph of a hand-drawn picture is acceptable.

You can earn 1 points of extra credit by submitting exactly 2 problems before 9am on 13 October 2021. Resubmissions void the extra credit.

1. [2] Show that there exists a constant c with the following property. Whenever G is a graph on n vertices and $\frac{1}{2}\binom{n}{2}$ edges, then there is a subset U of vertices such that $|e(U) - \frac{1}{2}\binom{|U|}{2}| \geq cn^{3/2}$.
2. [2] Prove that there is an absolute constant $c > 0$ with the following property. Let A be an n -by- n matrix with pairwise distinct real entries. Then there is a permutation of the rows of A so that no column in the permuted matrix contains an increasing subsequence of length at least $c\sqrt{n}$.
3. [2+(1 extra credit)]
 - (a) Let \mathcal{S} be a set of binary words of finite length and assume that no word in \mathcal{S} is a prefix of another. Let N_ℓ be the number of words of length ℓ in \mathcal{S} . Prove that $\sum_\ell N_\ell 2^{-\ell} \leq 1$.
 - (b) Let $w_1 w_2$ denote the concatenation of words w_1 and w_2 . Suppose that \mathcal{S} is a set of binary words of finite length such that for every word w there is at most one way of writing it as $w = w_1 \dots w_k$ for some $k \in \mathbb{Z}_+$ and some $w_i \in \mathcal{S}$. Prove that the inequality $\sum_\ell N_\ell 2^{-\ell} \leq 1$ continues to hold.
4. [1] Let A be a random subset of $[n]$ such that $\Pr[k \in A] = p$ and the events $k \in A$ are independent. Let E denote the event that A contains a 4-term arithmetic progression.

*This homework is from <http://www.borisbukh.org/DiscreteMath21/hw4.pdf>.

Find a function $t(n)$ such that

- if $p = o(t(n))$, then $\Pr[E] \rightarrow 0$; and
- if $p = \omega(t(n))$, then $\Pr[E] \rightarrow 1$.

5. [2] Write down an explicit positive constant c such that the following holds for all natural numbers n . Suppose a_1, \dots, a_n are n real numbers satisfying $\sum_{i=1}^n a_i^2 = 1$, and $\epsilon_1, \dots, \epsilon_n \in \{-1, +1\}$ are uniformly and independently chosen signs. Then

$$\Pr \left[\left| \sum_{i=1}^n \epsilon_i a_i \right| \leq 1 \right] \geq c.$$